## MHS Health Wisconsin Calendar Year 2020 HEDIS® ratings:

The Healthcare Effectiveness Data and Information Set (HEDIS<sup>®</sup>) is a tool that helps health plans measure the quality of care their members receive. MHS Health uses HEDIS<sup>®</sup> data to identify areas for improvement and monitor progress. The chart below shows our progress on HEDIS measures and areas that need improvement, based on our Calendar Year (CY) 2020 HEDIS<sup>®</sup> scores.

Measure	CY2020 Results	Percentile Ranking	CY2020 Goal NCQA Quality Compass 75 <sup>th</sup> Percentile	CY2019 Results
<b>Goal Met or Exceede</b> Adherence to Antipsychotic Medications for Individuals with Schizophrenia	d 72.08	75 <sup>th</sup>	68.90	71.46
	53.80	90 <sup>th</sup>	45.61	48.35
Antidepressant Medication Management Continuation		75 <sup>th</sup>		
Appropriate Testing for Pharyngitis	84.93		82.20	82.25
Pharmacotherapy Management of COPD Exacerbation: Corticosteroid	78.21	75 <sup>th</sup>	75.47	78.37
Pharmacotherapy Management of COPD Exacerbation: Bronchodilator	90.38	75 <sup>th</sup>	88.11	88.46
Statin Therapy for Patients with Cardiovascular Disease - Received Statin Therapy	85.79	90 <sup>th</sup>	82.90	84.35
Statin Therapy for Patients with Cardiovascular Disease - Statin Adherence	81.60	90 <sup>th</sup>	76.98	87.10
Statin Therapy for Patients with Diabetes - Received Statin Therapy	69.82	75 <sup>th</sup>	69.33	67.59
Statin Therapy for Patients with Diabetes - Statin Adherence	84.59	95 <sup>th</sup>	73.43	76.28
Use of Imaging Studies for Low Back Pain	81.28	75 <sup>th</sup>	79.53	80.00
On Track with National Av	erages			
Avoidance of Antibiotic Treatment in Acute Bronchitis/Bronchiolitis	54.25	50 <sup>th</sup>	61.90	51.11
Breast Cancer Screening	56.89	67 <sup>th</sup>	58.70	65.00
Cervical Cancer Screening	61.17	50 <sup>th</sup>	63.66	65.43
Childhood Immunization Status (Combo 10)	41.85	50 <sup>th</sup>	42.34	34.80
Chlamydia Screening in Women	55.08	50 <sup>5h</sup>	61.75	55.18
Comprehensive Diabetes Care: BP Control <140/90	64.96	67 <sup>th</sup>	65.69	67.15
Controlling High Blood Pressure	58.88	50 <sup>th</sup>	62.53	64.76
Follow-Up After Emergency Department Visit for Alcohol or Other Drug Abuse or Dependence: 7 days	15.12	50 <sup>th</sup>	17.66	15.63
Follow-Up After Emergency Department Visit for Mental Illness: 7 days	44.44	50 <sup>th</sup>	49.49	50.73
Follow-Up After Hospitalization for Mental Illness: 7 days	47.75	67 <sup>th</sup>	49.49	46.52
Initiation and Engagement of Alcohol and Other Drug Dependence Treatment: Engagement	14.72	50 <sup>th</sup>	17.66	15.79
Prenatal Care	88.08	50 <sup>th</sup>	89.29	88.08
Needs Improvement				
Annual Dental Visits	25.27	10 <sup>th</sup>	52.46	37.64
Appropriate Treatment for Upper Respiratory Infection	88.15	33 <sup>rd</sup>	92.11	90.42
Asthma Medication Ratio	59.21	10 <sup>th</sup>	70.67	54.62
Comprehensive Diabetes Care: Eye Exam	50.61	33 <sup>rd</sup>	57.91	66.42
Comprehensive Diabetes Care: HbA1c Control (<8.0%)	45.26	33 <sup>rd</sup>	51.34	52.31
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who are Using Antipsychotic	75.42	33 <sup>rd</sup>	79.70	78.05
Medications Follow-Up for Children on ADHD Medication -	38.89	10 <sup>th</sup>	62.44	36.84
Continuation Phase Immunizations for Adolescents:	29.99	10 <sup>th</sup>	43.55	33.60
Combination 2 Metabolic Monitoring for Children and Adolescents on Antipsychotics	17.05	<5 <sup>th</sup>	36.84	21.62
Postpartum Care	71.05	<5 10 <sup>th</sup>	79.56	73.48

## MHS Health Wisconsin Calendar Year 2020 HEDIS® ratings:

The Healthcare Effectiveness Data and Information Set (HEDIS<sup>®</sup>) is a tool that helps health plans measure the quality of care their members receive. MHS Health uses HEDIS<sup>®</sup> data to identify areas for improvement and monitor progress. The chart below shows our progress on HEDIS measures and areas that need improvement, based on our Calendar Year (CY) 2020 HEDIS<sup>®</sup> scores.

Measure	CY2020 Results	Percentile Ranking	CY2020 Goal NCQA Quality Compass 75 <sup>th</sup> Percentile	CY2019 Results
Risk of Continued Opioid Use - 31 day (lower is better)	7.03	10 <sup>th</sup>	2.43	6.42
Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics	50.00	10 <sup>th</sup>	69.50	56.41
Use of Opioids at High Dosage (lower rate is better)	20.88	5 <sup>th</sup>	2.38	15.94
Use of Opioids from Multiple Providers - Multipe Prescribers and Multiple Pharmacies (lower rate is better)	2.76	25 <sup>th</sup>	0.93	4.09
Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents: BMI Percentile	76.31	33 <sup>rd</sup>	82.73	65.70