

## **Clinical Policy: Factor IX (Human, Recombinant)**

Reference Number: CP.PHAR.218

Effective Date: 05.01.16

Last Review Date: 02.19

Line of Business: Medicaid, HIM-Medical Benefit

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

The following are factor IX products requiring prior authorization: human – AlphaNine SD<sup>®</sup>, Mononine<sup>®</sup>; recombinant – Alprolix<sup>®</sup>, BeneFIX<sup>®</sup>, Idelvion<sup>®</sup>, Ixinity<sup>®</sup>, Rebinyn<sup>®</sup>, Rixubis<sup>®</sup>.

### **FDA Approved Indication(s)**

Factor IX products are indicated for patients with hemophilia B (congenital factor IX deficiency or Christmas disease) for the following uses:

- Prevention and control of bleeding (on-demand treatment)
  - Adults and children: AlphaNine SD, Alprolix, BeneFIX, Idelvion, Ixinity (≥ 12 years), Mononine, Rebinyn, and Rixubis
- Perioperative management of bleeding
  - Adults and children: Alprolix, BeneFIX, Idelvion, Ixinity (≥ 12 years), Rebinyn, and Rixubis
- Routine prophylaxis to reduce the frequency of bleeding episodes
  - Adults and children: Alprolix, Idelvion, and Rixubis

Limitation(s) of use:

- AlphaNine SD, BeneFIX, and Mononine contain low, non-therapeutic levels of factors II, VII, and X, and, therefore, are not indicated for the treatment of factor II, VII or X deficiencies. They are also not indicated for the reversal of coumarin anticoagulant-induced hemorrhage, nor in the treatment of hemophilia A patients with inhibitors to factor VIII.
- BeneFIX and Mononine are also not indicated in a hemorrhagic state caused by hepatitis-induced lack of production of liver dependent coagulation factors.
- Alprolix, Idelvion, Ixinity, Rebinyn, and Rixubis are not indicated for induction of immune tolerance in patients with hemophilia B.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that AlphaNine SD, Alprolix, BeneFIX, Idelvion, Ixinity, Mononine, Rebinyn, and Rixubis are **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Congenital Hemophilia B (must meet all):**

1. Diagnosis of congenital hemophilia B (factor IX deficiency);

**CLINICAL POLICY****Factor IX (Human, Recombinant)**

2. Prescribed by or in consultation with a hematologist;
3. Age  $\geq$  17 years (AlphaNine only) or  $\geq$  12 years (Ixinity only);
4. Request is for one of the following uses (a, b, or c):
  - a. Control and prevention of bleeding episodes;
  - b. Perioperative management;
  - c. Routine prophylaxis to prevent or reduce the frequency of bleeding episodes (Alprolix, Idelvion, or Rixubis only);
5. Dose does not exceed the FDA approved maximum recommended dose for the relevant indication.

**Approval duration: 3 months (surgical/acute bleeding) or 6 months (prophylaxis)**

**B. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid and HIM-Medical Benefit.

**II. Continued Therapy****A. Congenital Hemophilia B (must meet all):**

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed the FDA approved maximum recommended dose for the relevant indication.

**Approval duration: 3 months (surgical/acute bleeding) or 6 months (prophylaxis)**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.  
**Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid and HIM-Medical Benefit.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PMN.53 for Medicaid and HIM-Medical Benefit or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

*Appendix B: Therapeutic Alternatives*

Not applicable

## CLINICAL POLICY

### Factor IX (Human, Recombinant)

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
  - All products except AlphaNine SD: known history of hypersensitivity reactions, including anaphylaxis, to the product or its excipients\*
    - \*Including mouse or hamster protein for BeneFix, Idelvion, Ixinity, Mononine, Rebinyn, and Rixubis
  - Rixubis: disseminated intravascular coagulation, signs of fibrinolysis
- Boxed warning(s): none reported

#### V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Factor IX, human (AlphaNine SD)	Control and prevention of bleeding episodes	<p>Minor episodes: 20-30 IU/kg IV twice daily</p> <p>Moderate episodes: 25-50 IU/kg IV twice daily</p> <p>Major episodes: 30-50 IU/kg IV twice daily for at least 3-5 days, followed by 20 IU/kg IV twice daily</p> <p>Surgery: 50-100 IU/kg IV twice daily before surgery, followed by the same regimen for 7-10 days thereafter</p>	<p>Bleeding episodes: 100 IU/kg/day</p> <p>Surgery: 200 IU/kg/day</p>
Factor IX, human (Mononine)	Control and prevention of bleeding episodes	<p>Minor episodes: 20-30 IU/kg IV every 24 hours</p> <p>Major trauma or surgery: 75 IU/kg IV every 18-30 hours</p>	<p>Minor episodes: 30 IU/kg/day</p> <p>Major trauma or surgery: 750 IU/kg/18 hours</p>
Factor IX, recombinant (Alprolix)	Control and prevention of bleeding episodes, perioperative management	<p>Minor and moderate episodes: 30-60 IU/dL/kg IV every 48 hours if there is further evidence of bleeding after the first dose</p> <p>Major episodes: 80-100 IU/dL/kg IV initially; consider a repeat dose after 6-10 hours and then every 24 hours for the first 3 days. May extend to dosing every 48 hours or longer after the first 3 days</p>	<p>Bleeding episodes: 100 IU/dL/kg/dose</p> <p>Surgery: 80 IU/dL/kg/dose</p>

**CLINICAL POLICY**  
**Factor IX (Human, Recombinant)**

Drug Name	Indication	Dosing Regimen	Maximum Dose
		<p>Minor surgery: 50-80 IU/dL/kg IV initially followed by every 24-48 hours until bleeding stops and healing is achieved</p> <p>Major surgery: 60-80 IU/dL/kg IV initially; consider a repeat dose after 6-10 hours and then every 24 hours for the first 3 days. May extend to dosing every 48 hours or longer after the first 3 days</p>	
	Routine prophylaxis	50 IU/dL/kg IV once weekly or 100 IU/dL/kg IV once every 10 days (start with 60 IU/kg once weekly for < 12 years)	100 IU/dL/kg/dose
Factor IX, recombinant (BeneFIX)	Control and prevention of bleeding episodes, perioperative management	<p>Minor episodes: 20-30 IU/dL/kg IV every 12-24 hours</p> <p>Moderate episodes: 25-50 IU/dL/kg IV every 12-24 hours</p> <p>Major episodes: 50-100 IU/dL/kg IV every 12-24 hours</p> <p>Surgery: 50-100 IU/dL/kg IV every 12-24 hours</p>	200 IU/dL/kg/day
Factor IX, recombinant (Idelvion)	Control and prevention of bleeding episodes, perioperative management	<p>Minor and moderate episodes: 30-60 IU/dL/kg IV every 48-72 hours</p> <p>Major episodes: 60-100 IU/dL/kg IV every 48-72 hours until bleeding stops and healing is achieved; maintenance dose is weekly</p> <p>Minor surgery: 50-80 IU/dL/kg IV every 48-72 hours until healing is achieved</p>	<p>Bleeding episodes: 100 IU/dL/kg/48 hours</p> <p>Surgery: 80 IU/dL/kg/48 hours</p>

**CLINICAL POLICY**  
**Factor IX (Human, Recombinant)**

Drug Name	Indication	Dosing Regimen	Maximum Dose
		Major surgery: 60-100 IU/dL/kg IV every 48-72 hours until bleeding stops and healing is achieved; maintenance dose is 1-2 times per week	
	Routine prophylaxis	<p>≥ 12 years of age: 25-40 IU/dL/kg IV every 7 days followed by 50-75 IU/dL/kg IV every 14 days once well-controlled</p> <p>&lt; 12 years of age: 40-55 IU/dL/kg IV every 7 days</p>	55 IU/dL/kg/week
Factor IX, recombinant (Ixinity)	Control and prevention of bleeding episodes, perioperative management	<p>Minor episodes: 30-60 IU/dL/kg IV every 24 hours</p> <p>Moderate episodes: 40-60 IU/dL/kg IV every 24 hours</p> <p>Major episodes: 60-100 IU/dL/kg IV every 12-24 hours</p> <p>Minor surgery: 50-80 IU/dL/kg IV pre-operatively followed by 30-80 IU/dL/kg every 24 hours</p> <p>Major surgery: 60-80 IU/dL/kg IV pre-operatively followed by 40-60 IU/dL/kg IV every 8-24 hours for 1-3 days or 30-50 IU/dL/kg IV every 8-24 hours for 4-6 days or 20-40 IU/dL/kg IV every 8-24 hours for 7-14 days</p>	<p>Bleeding episodes: 102 IU/dL/kg/dose</p> <p>Surgery: 81.6 IU/dL/kg/dose</p>
Factor IX, recombinant (Rixubis)	Control and prevention of bleeding episodes, perioperative management	<p>Minor episodes: 20-30 IU/dL/kg IV every 12-24 hours until healing is achieved</p> <p>Moderate episodes: 25-50 IU/dL/kg IV every 12-24 hours until bleeding stops and healing is achieved</p>	100 IU/dL/kg/dose

**CLINICAL POLICY**  
**Factor IX (Human, Recombinant)**

Drug Name	Indication	Dosing Regimen	Maximum Dose
		Major episodes: 50-100 IU/dL/kg IV every 12-24 hours until bleeding stops and healing is achieved  Minor surgery: 30-60 IU/dL/kg IV every 24 hours until healing is achieved  Major surgery: 80-100 IU/dL/kg IV every 8-24 hours until bleeding stops and healing is achieved	
	Routine prophylaxis	≥ 12 years of age: 40-60 IU/dL/kg IV twice weekly < 12 years of age: 60-80 IU/dL/kg IV twice weekly	80 IU/dL/kg/dose
Factor IX, recombinant, glycopegylated (Rebinyn)	On-demand treatment and control of bleeding episodes	40 IU/kg body weight for minor and moderate bleeds, and 80 IU/kg body weight for major bleeds. Additional doses of 40 IU/kg can be given	80 IU/kg/dose
	Perioperative management of bleeding	Pre-operative dose of 40 IU/kg body weight for minor surgery, and 80 IU/kg body weight for major surgery. As clinically needed for the perioperative management of bleeding, repeated doses of 40 IU/kg (in 1-3 day intervals) within the first week after major surgery may be administered. Frequency may be extended to once weekly after the first week until bleeding stops and healing is achieved	80 IU/kg pre-operatively; 40 IU/kg/dose after surgery

**VI. Product Availability**

Drug Name	Availability
Factor IX, human (AlphaNine SD)	Vial: 500, 1,000, 1,500 IU
Factor IX, human (Mononine)	Vial: 500, 1,000 IU
Factor IX, recombinant (Alprolix)	Vial: 250, 500, 1,000, 2,000, 3,000, 4,000 IU
Factor IX, recombinant (BeneFIX)	Vial: 250, 500, 1,000, 2,000, 3,000 IU
Factor IX, recombinant (Idelvion)	Vial: 250, 500, 1,000, 2,000 IU

**CLINICAL POLICY**  
**Factor IX (Human, Recombinant)**

Drug Name	Availability
Factor IX, recombinant (Ixinity)	Vial: 250, 500, 1,000, 1,500, 2,000, 3,000 IU
Factor IX, recombinant (Rixubis)	Vial: 250, 500, 1,000, 2,000, 3,000 IU
Factor IX, recombinant, glycopegylated (Rebinyn)	Vial: 500, 1,000, 1,500 IU

**VII. References**

1. Alphanine SD Prescribing Information. Los Angeles, CA: Grifols Biologicals, Inc.; March 2017. Available at: [www.alphaninesd.com](http://www.alphaninesd.com). Accessed November 8, 2018.
2. Alprolix Prescribing Information. Cambridge, MA: Biogen Idec, Inc.; June 2018. Available at: [www.alprolix.com](http://www.alprolix.com). Accessed November 8, 2018.
3. BeneFix Prescribing Information. Philadelphia, PA: Wyeth Pharmaceuticals, Inc.; June 2017. Available at: [www.benefix.com](http://www.benefix.com). Accessed November 8, 2018.
4. Idelvion Prescribing Information. Kankakee, IL: CSL Behring LLC; May 2018. Available at: [www.idelvion.com](http://www.idelvion.com). Accessed November 8, 2018.
5. Ixinity Prescribing Information. Berwyn, PA: Aptevo BioTherapeutics LLC; April 2018. Available at: [www.ixinity.com](http://www.ixinity.com). Accessed November 8, 2018.
6. Mononine Prescribing Information. Kankakee, IL: ZLB Behring, LLC; April 2016. Available at: [www.http://labeling.cslbehring.com/PI/US/Mononine/EN/Mononine-Prescribing-Information.pdf](http://www.http://labeling.cslbehring.com/PI/US/Mononine/EN/Mononine-Prescribing-Information.pdf). Accessed November 8, 2018.
7. Rixubis Prescribing Information. Westlake Village, CA: Baxalta US Inc.; March 2016. Available at: <http://www.rixubis.com>. Accessed November 8, 2018.
8. Srivastava A, Brewer AK, Mauser-Bunschoten EP, et al. Guidelines for the management of hemophilia. *Haemophilia*. Jan 2013; 19(1): e1-47.
9. Medical and Scientific Advisory Council (MASAC) of the National Hemophilia Foundation (NHF): Database of treatment guidelines. Available at <https://www.hemophilia.org/Researchers-Healthcare-Providers/Medical-and-Scientific-Advisory-Council-MASAC/MASAC-Recommendations>. Accessed September 26, 2018.
10. Rebinyn Prescribing Information. Plainsboro, NJ: Novo Nordisk; May 2017. Available at: [www.rebinyn.com](http://www.rebinyn.com). Accessed November 8, 2018.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J7194	Factor IX complex, per IU
J7195	Injection, factor IX (antihemophilic factor, recombinant) per IU, not otherwise specified
J7200	Injection, factor IX, (antihemophilic factor, recombinant), Rixubis, per IU
J7201	Injection, factor IX, FC fusion protein (recombinant), per IU
J7202	Injection, factor IX, albumin fusion protein, (recombinant), Idelvion, per IU.

**CLINICAL POLICY**  
**Factor IX (Human, Recombinant)**

Reviews, Revisions, and Approvals	Date	P&T Approval Date
<p>Policy split from CP.PHAR.12.Blood Factors and converted to new template. Removed requests for documentation.            Added age requirement per PI for Ixinity.            Under initial criteria, removed requirement for “history of 2 or more joint bleeds.” Delineated Alprolix and Rixubis for prophylaxis per Pis.            Approval period for non-prophylactic use is edited to provide 3 months on initial approval and one 3-month re-auth; approval period for prophylactic use is retained at 6 months initial/6 months continuing therapy. Removed denial based on inhibitor titer of <math>\geq 5</math> BU/mL as Pis do not specify a limit.            Reviewed by specialist.</p>	04.01.16	05.16
<p>Safety information removed. Wording for uses of all blood factor products made consistent across all policies. Added indication for Alprolix and Rixubis for routine prophylaxis. Approval periods across all blood factor policies made consistent. Efficacy statement added to renewal criteria.            Hemophilias are specified as “congenital” versus “acquired” across blood factor policies where indicated. Reviewed by specialist-hematology/internal medicine.</p>	04.01.17	05.17
<p>1Q18 annual review:            - Converted to new template            - Added Idelvion to the policy under the same coverage criteria as the other recombinant factor IX agents.            - Specified routine prophylaxis indication is only for certain agents, per package labeling for those agents.            - Added age limit for AlphaNine per package labeling            - References reviewed and updated.</p>	11.28.17	02.18
<p>1Q 2019 annual review: added HIM-Medical Benefit; no significant changes; added Rebinyn; references reviewed and updated.</p>	11.08.18	02.19

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health



## CLINICAL POLICY

### Factor IX (Human, Recombinant)

plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members,** when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed,

## CLINICAL POLICY

### Factor IX (Human, Recombinant)

displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.